

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355822973>

Studies in Canarium urceus (Linné, 1758) Part 5: a New Species from the Northern Pacific Ocean (Gastropoda: Neostromboidae: Strombidae)

Article · November 2021

DOI: 10.54173/F534282

CITATIONS

0

READS

45

2 authors:



Stephen J Maxwell

James Cook University

41 PUBLICATIONS 59 CITATIONS

[SEE PROFILE](#)



Aart Marinus Dekkers

Independent researcher; CPA at Dekkers Accountants

60 PUBLICATIONS 69 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Miscellaneous [View project](#)



New species descriptions in Mitridae [View project](#)

Studies in *Canarium urceus* (Linné, 1758) Part 5: a New Species from the Northern Pacific Ocean (Gastropoda: Neostromboidae: Strombidae)

Stephen J. Maxwell¹ and Aart M. Dekkers²

¹ College of Science and Engineering, James Cook University, Cairns Qld 4870, Australia.

stephen.maxwell@my.jcu.edu.au

² Oasestraat 79, 1448 NR Purmerend, The Netherlands.

aart.dekkers@wxs.nl

ABSTRACT This part of the *Canarium (Canarium) urceus* (Linné, 1758) revision after Abbott (1960) revision examines the northwest Pacific specimens, which are currently synonymised under *Canarium (Canarium) urceus* (Linné, 1758). *Canarium (Canarium) nipponium* n. sp. is known from the Ryukyuan Subprovince, Mariana Islands and Palau. The new species differs from others in lacking the fine sculpture on the early whorls and triangulate form. Given the propensity for species of Strombidae to disperse widely, to show a high degree of variability across their ranges, and to hybridise, it is expected that traits of this new species will be present in populations of neighbouring species, particularly within the *Canarium (Canarium)* complex.

KEYWORDS *Canarium*, *C. urceus*, *C. nipponium*, Japan, Mariana Islands, Palau, *Strombus*, Ryukyuan Subprovince

INTRODUCTION

The *Canarium (Canarium) urceus* (Linné, 1758) complex (Abbott 1960) has recently been significantly revised, with the identification of many new species within bioregions of the Indo-Australian Archipelago realm (Maxwell *et al.* 2020a; 2020b; Dekkers and Maxwell 2020). There is a sound basis for the recognition of species based on biogeographic boundaries based on the phylogenetic history of a diverse set of marine organisms within the Indo-Australian Archipelago realm (Santini and Winterbottom 2002; Spalding *et al.* 2007; Carpenter *et al.* 2011; Borsa *et al.* 2016; Kulbicki *et al.* 2013; Veron *et al.* 2015; Yang *et al.* 2016; Wainwright *et al.* 2018). The distribution of *Canarium* examined in this study falls within the Ryukyuan Subprovince and the Mariana Chain of Islands, which is considered outside and to the north of the Indo-Australian

Archipelago realm (Petuch and Berschauer 2020; Figure 1).

ABBREVIATIONS

- AMD Aart M. Dekkers Collection, Purmerend, The Netherlands.
SMC Stephen Maxwell Collection, Cairns, Queensland, Australia.

METHODS

The rarely observed species described in this paper is based on two specimens, one from the Hahajima Retto, Ogasawara Islands, and one collected from Ryukyu (Figure 1). The Ryukyu specimen was obtained from Thora Whitehead, Brisbane, Australia, and was collected prior to the 1970s. The Hahajima Retto, Ogasawara Islands specimen was obtained from a dealer, and was collected in the 1950s. The species is also illustrated in the molluscan literature of

Japan (Hirase 1936, 1954; Kira 1959). While the number of physical specimens at our disposal is limited, the morphological similarity of the material from two sources, the illustrations of it in literature (under the "*urceus*" name), and the morphological distinctiveness from other *Canarium* (*Canarium*) justifies their recognition.

The decision to use the rank of species or subspecies is based on Maxwell and Dekkers (2019a, b) and Maxwell *et al.* (2021). Under this system, the subspecies rank should be restricted to those taxa where there are no other forms of discrimination other than phenetic differences in genetic sequences.

SYSTEMATIC PART

| | |
|-------------|---|
| Superfamily | Stromoidea Rafinesque, 1815 |
| Epifamily | Neostromboidae Maxwell, Dekkers, Rymer & Congdon, 2019 |
| Family | Strombidae Rafinesque, 1815 |
| Subfamily | Neostrombinae Maxwell & Rymer, 2021 |
| Tribe | Neostrombini Liverani, Dekkers & Maxwell, 2021 |
| Genus | <i>Canarium</i> Schumacher, 1817 |
| Subgenus | <i>Canarium</i> Schumacher, 1817 (Type species – <i>Canarium</i> <i>ustulatum</i> Schumacher, 1817 = <i>Canarium (urceus) urceus</i> (Linné, 1758)) |

Canarium (Canarium) nipponium Maxwell &
Dekkers, new species
(Figure 1)

Synonymy.

1936 *Strombus ustulatus* Schumacher –
Hirase, pl. 86, fig. 11, and Hirase
1954, pl. 86, fig. 11.

1959 *Strombus ustulatum* Linné – Kira, p.
35, pl. 15, fig. 4.

Diagnosis. Solid shells with triangular body whorl. The shoulders of the whorls are angulated and the early whorls lack fine sculpture. The posterior aperture is restricted. Body whorl covered with many very narrow spiral lines.

Description. The shell is heavy and solid. The shape is ovate to fusiform with an extended anterior sinus (broken in holotype) that is not dorsally exposed. The stromboid sinus is acute and axially angled in the direction of shell growth, and has a thin protruding flange that is quadrate and typically dark stained with black to brown both externally and internally. The ventral shoulder has reduced blunt knobs that continue to the dorsum with three well defined blunt shoulder knobs, and one greatly reduced knob prior to the formation of the outer lip. There are fine axial incised lines on the lower third of the body whorl both dorsally and ventrally. Early teleoconch whorls are rounded and smooth, with a distinctive subsutural ramp. Later whorls become more angulate at the shoulder, developing axially elongated shoulder knobs, with fine well-spaced spiral incised lines that do not extend to the body whorl. The spire often has a purplish tint. The aperture is lanceolate, being restricted at the posterior end, white, with very thin brownish lines entering the aperture on the inside of the lip. The columella is well formed and uniform along the aperture, having a yellowish hue. The other lip is thickened and somewhat quadrate posteriorly, joining at the shoulder of the body whorl, becoming thinner and sharp-edged anteriorly.

Type Material. Holotype – Ryukyu, Japan 37 mm MNHN-IM-2000-35893; Paratype 1 – Hahajima, Ogasawara Islands, Japan, (SMC 19g. 001).

Type Locality. We designate Ryukyu, Japan as the type locality.

Etymology. Named after the country of the type population, Japan.

Locality Records. *Literary Records:* Japan – Ryūkyū (Hirase 1936, 1954); Yaka Beach, Ishikawa (Abbott 1960). *Material Examined:* Japan – Hahajima, Ogasawara Islands, Japan (SMC x 1); Ryūkyū (SMC x 1). Palau – Palau Lagoon (AMD x 3).

Comparison and Remarks. *Canarium* (*C.*) *nippонium* n. sp. is similar in form to *C.* (*C.*) *darwinensis* Maxwell and Dekkers, 2021 and *C.* (*C.*) *manintveldi* (Dekkers and Maxwell, 2020) but lacks the fine sculpture on the early whorls of that species. *C.* (*C.*) *incisum* (Wood, 1828) differs in having a uniform orange aperture, a more triangulate shape. *C.* (*C.*) *urceus* (Linné, 1758) differs in form, having a more angulate shoulder on the body whorl, and the posterior aperture not restricted as it is in *C.* (*C.*) *nippонium* n. sp.. *Canarium* (*C.*) *anatellum* (Duclos, 1844) and *C.* (*C.*) *esculentum* (Maxwell, Rymer, Congdon & Dekkers, 2020b) lack the angulated whorls and shouldered aperture of the new species. Material from online sources (www.stromboidea.de) and literary references, indicate that *C.* (*C.*) *nippонium* n. sp. is presently restricted to the Ryukyuan Subprovince of Japan and the Mariana Islands south to Palau.

DISCUSSION

There is growing evidence that the *C.* (*C.*) *urceus* (Linné, 1758) complex (Abbott 1960), despite variability in form and sharing of similar phenotypic traits (e.g. random presence of a black columella in some individuals), are also evolving in divergent directions, often within defined biogeographic provinces and zones. These recurrent morphological themes can be

explained in three ways: first, there is a lag between the time of genetic divergence and evidence of this split in the morphology of organisms (Baum 2008); second, within a population, there are enough carriers of a mutation that drift and selection maintain allelic frequency without leading to either extinction or fixation (Baum 2008); or third, there is enough genetic inflow from populations outside the particular bioregion that would enable recurrent introduction within a population of a particular morphological trait to sustain its ongoing recurrence (Delgado *et al.* 2006; Paris *et al.* 2006, 2008).

REFERENCES

- Abbott, R.T. 1960.** The genus *Strombus* in the Indo-Pacific. Indo-Pacific Mollusca 1(2):35-146.
- Baum, D. 2008.** Reading a Phylogenetic Tree: The Meaning of Monophyletic Groups. Nature Education 1(1):190.
- Borsa, P., K.-N. Shen, I.S. Arlyza, & T.B. Hoareau. 2016.** Multiple cryptic species in the blue-spotted maskray (Myliobatoidei: Dasyatidae: *Neotrygon* spp.): An update. Espèces cryptiques multiples chez la pastenague masquée à points bleus (Myliobatoidei: Dasyatidae: *Neotrygon* spp.): actualisation. Comptes Rendus Biologies 339(9-10):417-426.
<https://doi.org/10.1016/j.crvi.2016.07.004>
- Carpenter, K.E., P.H. Barber, E.D. Crandall, M.C.A. Ablan-Lagman, G.N. Ambariyanto, Mahardika, B.M. Manjaji-Matsumoto, M.A. Juinio-Menez, M.D. Santos, C.J. Starger & A.H.A. Toha. 2011.** Comparative phylogeography of the Coral Triangle and Implications for Marine Management. Journal of Marine Biology 2011:396982.
<https://doi.org/10.1155/2011/396982>

- Dekkers, A.M. & S.J. Maxwell.** 2020. Studies in *Canarium urceus* (Linnaeus, 1758) Part 3: new species from the western Pacific (Gastropoda: Neostromboidae: Strombidae). *The Festivus* 52(4):345-358. <https://doi.org/10.54173/F524345>
- Delgado, G.A., R.A. Glazer, D. Hawtof, D. Aldana Aranda, L.A. Rodríguez-Gil & A. de Jesús-Navarrete.** 2006. Do queen conch (*Strombus gigas*) larvae recruiting to the Florida Keys originate from upstream sources? Evidence from plankton and drifter studies. In Grober-Dunsmore, R., and B.D. Keller, eds., Caribbean connectivity: Implications for marine protected area management. Proceedings of a Special Symposium, 9-11 November 2006, 59th Annual Meeting of the Gulf and Caribbean Fisheries Institute, Belize City, Belize. pp. 29-41.
- Duclos, P.L. 1843-1844.** *Strombus*, pls. 1-30. In Chenu, J.C., 1842-1853, Illustrations conchyliologiques: ou description et figures de toutes les coquilles connues vivante et fossiles classes suivant le système de Lamarck modifié d'après les progress de la science, et comprenant les genres nouveaux et les espèces récemment découvertes. Fortin, Masson, Paris.
- Hirase, S. 1936.** A Collection of Japanese Shells with Illustrations in Natural Colours (5th edn). M. Sanshodo, Tokyo.
- Hirase, S. 1954.** An Illustrated Handbook or Shells in Natural Colour from the Japanese Islands and Adjacent Territories (Revised by I. Taki). Maruzen Co., Tokyo.
- Kira, T. 1959.** Coloured Illustrations of the Shells of Japan. Hoikusha Publishing Co., Osaka.
- Kulbicki, M., V. Parravicini, D.R. Bellwood, E. Arias-González, P. Chabanet, S.R. Floeter, A. Friedlander, J. McPherson, R.E. Myers, L. Vigliola & D. Mouillot.** 2013. Global Biogeography of Reef Fishes: A Hierarchical Quantitative Delineation of Regions. *PLoS One* 8(2):e18147. <https://doi.org/10.1371/journal.pone.0081847>
- Linné, C. 1758.** *Systema Naturae per Regna Tria Naturae Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis*, volume 1, 10th edn. Reformata. Holmiae, Laurentii Salvii. 824 pp. <https://doi.org/10.5962/bhl.title.542>
- Liverani, V., A.M. Dekkers & S.J. Maxwell.** 2021. Resolving phylogenetic and classical nomenclature: A Revision of *Canarium* Schumacher, 1817 (Mollusca, Neostromboidae, Strombidae). *The Festivus* 53(1):26-43. <https://doi.org/10.54173/F531026>
- Maxwell, S.J. & A.M. Dekkers.** 2019. A new name for *Altivasum typicum* Hedley, 1916 fide Dekkers and Maxwell, 2018 and the description of *Altivasum clarksoni nov. sp.*. *The Festivus* 51(2):171-176. <https://doi.org/10.54173/F513171>
- Maxwell, S.J. & A.M. Dekkers.** 2021. Studies in *Canarium urceus* (Linné, 1758) Part 4: *Canarium (Canarium) orrae* (Abbott, 1960) (Gastropoda: Neostromboidae: Strombidae) and a new species from the Northern Territory, Australia. *The Festivus* 53(4):270-281. <https://doi.org/10.54173/F534270>
- Maxwell, S.J. & T.L. Rymer.** 2021. Are the ICZN and PhyloCode that incompatible? A summary of the shifts in Stromboidean taxonomy and the definition of two new subfamilies in Strombidae (Mollusca, Neostromboidae). *The Festivus* 53(1):44-51. <https://doi.org/10.54173/F531044>
- Maxwell, S.J., A.V. Bordon, T.L. Rymer & B.C. Congdon.** 2019a. The birth of a species and the validity of hybrid

- nomenclature demonstrated with a revision of hybrid taxa within Strombidae (Neostromboidae). Proceedings of the Biological Society of Washington 132(1):119-130.
<https://doi.org/10.2988/19-00007>
- Maxwell, S.J., A.M. Dekkers, T.L. Rymer & B.C. Congdon. 2019b.** Recognising and defining a new crown clade within STROMBOIDEA Rafinesque, 1815 (MOLLUSCA, GASTROPODA). ZooKeys 867:1-7.
<https://doi.org/10.3897/zookeys.867.34381>
- Maxwell, S.J., T.L. Rymer, M.K. Rowell, L.C. Hernandez Duran, D.P. Berschauer, M. Underdown, E.J. & A.M. Dekkers. 2021.** Defining and Bringing Relevance of Meaning to Species Group-Level Taxa. Proceedings of the Biological Society of Washington 134:27-28.
<https://doi.org/10.2988/006-324X-134.1.27-28>
- Maxwell, S.J., T.L. Rymer & A.M. Dekkers. 2020a.** *Canarium urceus* (Linné, 1758) studies Part1: The Recircumscription of *Strombus urceus* Linné, 1758 (Neostromboidae: Strombidae). The Festivus 52(2):113-127.
<https://doi.org/10.54173/F522113>
- Maxwell, S.J., T.L. Rymer, B.C. Congdon & A.M. Dekkers. 2020b.** Studies in *Canarium urceus* (Linné, 1758) Part 2: *Strombus anatellus* Duclos, 1844, *Strombus crassilabrum* Anton, 1839, *Strombus incisus* Wood, 1828 and *Strombus ustulatus* form *laevis* Dodge, 1946 (Neostromboidae: Strombidae). The Festivus 52(4):335-344.
<https://doi.org/10.54173/F524335>
- Paris, C.B., M. Perez Perez, J. Kool & D. Aldana Aranda. 2006.** Segregation of queen conch, *Strombus gigas*, populations from the Yucatan Peninsula, Mexico. In Grober-Dunsmore, R., and B.D. Keller, eds., Caribbean connectivity: Implications for marine protected area management. Proceedings of a Special Symposium, 9-11 November 2006, 59th Annual Meeting of the Gulf and Caribbean Fisheries Institute, Belize City, Belize. pp. 71-88.
- Paris, C.B., D. Aldana Aranda, M. Perez Perez & J. Kool. 2008.** Connectivity of queen conch, *Strombus gigas*, populations from Mexico. Proceedings of the 11th International Coral Reef Symposium, Ft. Lauderdale, Florida, 7-11 July 2008, Session number 14, p. 439-443.
- Petuch, E.J. & D.P. Berschauer. 2020.** Tropical Marine Mollusks: An illustrated Biogeographical Guide. CRC Press, Boca Raton. 357 pp.
<https://doi.org/10.1201/9781003120070>
- Rafinesque, C.S. 1815.** Analyse de la Nature, ou tableau de l'Univers et des Corps Organisés. L'Imprimerie de Jean Barravecchia, Palermo. 224 pp.
<https://doi.org/10.5962/bhl.title.106607>
- Santini, F. & R. Winterbottom. 2002.** Historical biogeography of Indo-Western Pacific coral reef biota; Is the Indonesian region a centre of origin? Journal of Biogeography 29:189-205.
<https://doi.org/10.1046/j.1365-2699.2002.00669.x>
- Schumacher, C.F. 1817.** Essai d'un Nouveau Système des Habitations des vers Testacés. Schultz, Copenhagen. 287 pp.
- Spalding, M.D., H.E. Fox, G.R. Allen, N. Davidson, Z.A. Ferdaña, M. Finlayson, B.S. Halpern, M.A. Jorge, A. Lombana, S.A. Lourie, K.D. Martin, E. McManus, J. Molnar, C.A. Recchia & J. Robertson. 2007.** Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. BioScience 57(7):573-583.
<https://doi.org/10.1641/B570707>

Veron, J., M. Stafford-Smith, L. DeVantier & E. Turak. 2015. Overview of distribution patterns of zooxanthellate Scleractinia. *Frontiers in Marine Science* 1:81.
<https://doi.org/10.3389/fmars.2014.00081>

Wainwright, B.J., I.S. Arlyza & S.A. Karl. 2018. Population genetics of the collector urchin, *Tripneustes gratilla*, in the Indonesian archipelago. *Marine Ecology* 39(6):e12530.
<https://doi.org/10.1111/maec.12530>

Wood, W. 1828. Supplement to the Index Testaceologicus; or A Catalogue of Shells,

British and Foreign. Richard Taylor, London. 59 pp.

<https://doi.org/10.5962/bhl.title.30661>

Yang, Y., N.C. Duke, F. Peng, J. Li, S. Yang, C. Zhong, R. Zhou & S. Shi. 2016. Ancient geographical barriers drive differentiation among *Sonneratia caseolaris* populations and recent divergence from *S. lanceolata*. *Frontiers in Plant Science* 7:1618.
<https://doi.org/10.3389/fpls.2016.01618>

Cite as: Maxwell, S.J. and A.M. Dekkers. 2021. Studies in *Canarium urceus* (Linné, 1758) Part 5: a New Species from the Northern Pacific Ocean (Gastropoda: Neostrombidae: Strombidae). *The Festivus* 53(4):282-287.
<https://doi.org/10.54173/F534282>




Figure 1. Distribution of *Canarium (Canarium) nipponium* n. sp.: A) Holotype – Ryukyu, Japan 37 mm MNHN-IM-2000-35893; B) Paratype 1 – Hahajima Retto, Ogasawara Islands, Japan, (SMC 19g. 001) (Image modified from Google Earth, accessed August 2021).